



Growth of Wheat and Na⁺/ K⁺ Ratio under Neutral and Alkaline Salts Stress

**Badar-Uz-Zaman^{1*}, Huda Khan², Muhammad Suhaib¹, Muhammad Ishaq³
Muhammad Arshadullah¹ and Imdad Ali Mahmood¹**

¹Land Resource Research Institute (LRR), National Agricultural Research Centre (NARC), Islamabad, Pakistan.

²Department of Biological Sciences, Karakoram International University, Gilgit Baltistan, Pakistan.

³Division of Soil and Water Conservation, Department of Soil Science, Pir Mehar Ali Shah University of Arid Agriculture (UAAR), Rawalpindi, Pakistan.

Authors' contributions

This work was carried out in collaboration among all authors. Author BUZ designed the study, wrote the protocol and conducted study with author HK. Authors MS and MA managed the literature searches. Author MI performed the statistical analysis. Author IAM provided chemical and germplasm for the experiment. Author BUZ wrote the final draft of the manuscript. All authors read and approved the final manuscript.

Article Information

DOI: 10.9734/ASRJ/2019/v2i430057

Editor(s):

(1) Dr. Alessandro Buccolieri, Department of Biological and Environmental Sciences and Technologies (DiSTeBA), Università del Salento, Italy.

Reviewers:

(1) Jagruti H. Sanghvi, India.

(2) I. Rashmi, ICAR-Indian Institute of Soil and Water Conservation, India.

Complete Peer review History: <http://www.sdiarticle4.com/review-history/53782>

Received 02 November 2019

Accepted 07 January 2020

Published 15 January 2020

Original Research Article

ABSTRACT

Saline soils contain multiple types of soluble salts. Role of alkaline salts (AS) may be different than neutral salts (NS) for crop growth. Usually these types of salinity frequently co-exist. Neutral salts generally induce osmotic stress and ion-induced injury to growing crop plants. Behavior of AS may be different due to high pH. This study was conducted to see the response of wheat growth and Na⁺/ K⁺ ratio under AS and NS application as nutrient solution study. Germinated disinfected seeds of wheat (Cv. Pak-13) and raised in sand. Ten days seedlings were transferred to pots containing standard nutrients solution, with the application of 30 and 60 mM of NS (NaCl, Na₂SO₄) and AS (NaHCO₃ and Na₂CO₃) separately in 1:1 ratio using complete randomize design in triplicates. The seedling growth period in salt solution comprised 33 days. Biomass and the selected inorganic ions were affected significantly (p< 0.01) under salt stress. The magnitude of loss of bio mass was 11

*Corresponding author: E-mail: badruar@yahoo.com, badruaar@yahoo.com;

percent higher in AS than that of NS application. Phosphorus and sulphur concentration were lower 10 and 7 percent with AS than NS respectively besides high Na^+/K^+ ratio and pH. Further such studies on salt tolerant crop varieties can be carried out to differentiate response under different types of salts.

Keywords: *Wheat; types of salt stress; bio-mass; phosphorus; sulphur; Na⁺/K⁺ ratio.*

1. INTRODUCTION

On earth, salts are 'the life line' of living organisms. When salts are dissolved in water, their function is dependent on the nature of the salt and the living recipient. Out of all living organisms, the plants categorized as glycophytes can only survive when salts in soils are available to the root system in a limited amount (usually $\text{ECe} < 4.0 \text{ dSm}^{-1}$). Low ECe is a 'salvation' for crop plants. The composition of salts determines most of the chemical processes surrounding the roots of a crop plant. When soil nutrients cross a threshold of salts, then it gives rise to soil salinity problems. Saline soils contain multiple types of soluble salt components and the compositions of soluble salts in saline soils are quite different among locations [1]. Saline soils are heterogenic in salt types. Soil salinization is an important agriculture-related problem, which frequently co-exists with salt and alkali stresses [2]. Soil salinity is composed of different types of salts, mainly neutral and alkaline salts. Neutral salts (NS) compose NaCl and Na_2SO_4 . Alkaline salts (AS) are consisted of NaHCO_3 and Na_2CO_3 . Salts in soil cause osmotic stress, nutritional effects, ion toxicity and physiological disorders [3] for growing crops. Salt stress exerts distinct effects on growth, ion balance, compatible solutes, and metabolism in old and young leaves [4]. Increasing salinity stress is antagonistic to fresh mass of plant [5]. Alkaline salts are understood as more destructive to plants at different growth stages than neutral salts [6]. This type of salt stress causes high pH stress [7], and can seriously affect mineral nutrition in the form of ion imbalances. This may also affect the oxygen supply capacity of the roots besides structure and function of root cells. Many saline soils are also alkaline due to the presence of sodium carbonates [8]. Alkaline salt stress is linked with high pH [9,10]. Neutral salts generally induce osmotic stress and ion-induced injury to crop plants [11]. Such salts cause harm to the plant mainly through osmotic stress and ion toxicity. Neutral and alkaline salts usually coexist in saline, saline sodic or sodic soils. The existence of such types of salts may affect crop growth separately or collectively, depending upon their origin. It is difficult to quantify such

types of salts in a heterogeneous system of soils, although their separate effect on plant growth can be studied in a hydroponics system. Neutral and alkaline salt stress may affect growth of wheat differentially and Na^+/K^+ ratio may be higher under neutral salt stress. Alkaline salt stress may cause stronger destructive effects on plants than neutral salt stress and the response of wheat may be variable.

2. MATERIALS AND METHODS

Treated seeds of *Triticum aestivum* (cv. Pak-13) with sodium hypochlorite (1% w/v) for 15 minutes [12] and then germinated in quartz sand, moist with distilled water. Foam-plugged the seedlings in lids of plastic pots containing 2.5 L of continuously aerated nutrient solution as constituted by Hoagland and Arnon [13]. After 12 days of sowing, applied neutral salts solution ($\text{NaCl}:\text{Na}_2\text{SO}_4=1:1$) @ 30 and 60 mM. Added separately alkaline salts solution ($\text{NaHCO}_3:\text{Na}_2\text{CO}_3=1:1$) @ 30 and 60 mM. Replaced the nutrient solution weekly. Light intensity remained at $450 \mu\text{mol m}^{-2} \text{S}^{-1}$. Adjusted photoperiod at 16 hd^{-1} light period and maintained temperature at $25 \pm 2^\circ\text{C}$. Unplugged the whole plants on the 32nd day of transplantation to pots. Rinsed the plants with distilled water, blotted with tissue paper and recorded their fresh mass. Measured root length and shoot height using a centimeter scale. Dried the plant samples at 65°C to constant mass. Recorded dry mass of each sample and digested samples in 2% nitric acid solution along with a few drops of perchloric acid. Determined phosphorus in the plant digested material as described by Jackson [14]. Analyzed sulphur in the digested material as given by Verma [15]. By flame photometry determined the concentrations of Na^+ and K^+ ions. Analyzed the data according to two factors complete randomized design and compared treatment means using LSD test by using the statistical software, [16].

3. RESULTS AND DISCUSSION

Seedlings of wheat (cv. Pak-13) responded significantly ($p \leq 0.01$) for growth and ion relations on the application of neutral and alkaline salts.

3.1 Effect of Types of Salt Stress on Bio Mass

Shoot fresh mass (SFM) decreased 4 and 7 percent on the application of neutral salts (NS) solution (NaCl: Na₂SO₄=1:1) @ 30 and 60 mM respectively. On the application of alkaline salts (AS) solution (NaHCO₃:Na₂CO₃=1:1) @ 30 and 60 mM, SFM decreased 12 and 20 percent respectively. Root fresh mass (RFM) decreased 5 and 10 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60mM RFM decreased 15 and 18 percent respectively. Shoot dry mass (SDM) decreased 9 and 12 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60 mM, SDM decreased 18 and 21 percent respectively. Root dry mass (RDM) decreased 5 and 8 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60mM, RDM decreased 13 and 16 percent respectively (Table 1).

Shoot height (SH) decreased 5 and 10 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS

solution @ 30 and 60 mM, SH decreased 15 and 20 percent respectively. Root length (RL) decreased 8 and 13 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60mM, RL decreased 18 and 22 percent respectively (Fig. 1).

Under the influence of both types of stresses, the magnitude of bio mass of shoot and root of wheat (cv. Pak-13) decreased. Alkaline salts affect pH of the nutrients supply to the roots. Such salts keep pH high, following restricted availability to macronutrients. Under osmotic effect of dissolved ions, the availability of potassium ion also remains limited especially in the presence of sodium ion. In this study the type of salts, from lower to higher concentrations, affected biomass of shoot and root system. Higher concentration of sodium ion caused deficiency of water in plant tissue as found by [5]. Sodium ion stress interferes biochemical processes [17] such as interruption of membranes, imbalance in nutrient ratio, damages the ability to detoxify reactive oxygen species and decreased photosynthetic activity. Also, lesser availability of water caused lesser flow of nutrients to the shoot system

Table 1. Impact of neutral salts (NS) and alkaline salts (AS) stresses on the bio mass (mg plant⁻¹) of wheat.

Salinity (mM)	Fresh mass		Dry mass	
	Shoot	Root	Shoot	Root
Control	826.5 a	461.5 a	41.3 a	23.1 a
NS (30)	793.4 b	438.4 b	37.6 b	21.9 b
NS (60)	760.4 c	415.4 c	36.4 c	21.2 c
AS (30)	727.3 d	392.3 d	33.9 d	20.1 d
AS (60)	661.1 e	378.5 e	32.7 e	19.4 e

Means sharing similar letter(s) for a parameter in a column do not differ significantly at $p < 0.01$; CV ($p < 0.01$) = 6.98 percent




Fig. 1. Effect of neutral salts (NS) and alkaline salts (AS) on shoot height and root length of wheat

from root [18]. Fresh mass of plant material consists of tissue besides water and other chemicals. As water maintains physico-chemical properties of a cell or an organ, therefore in fresh mass consideration water contents of an organ cannot be ignored [19]. Dry mass of shoot and root decreased under both types of applied salts. When the relations and metabolic activities of macro and micro nutrients are on positive side of metabolism, then tissue developments in growing plants continues and this positive effect is reflected as dry mass of a plant organ.

3.2 Effect of Types of Salt Stress on Sulphur and Phosphorus Concentration in Wheat

Sulphur in shoot decreased 5 and 10 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60 mM, sulphur decreased in shoot 15 and 20 percent respectively. Sulphur in root decreased 6 and 14 percent on the application of NS solution @ 30 and 60mM respectively. On the application of AS solution @ 30 and 60mM, sulphur decreased in root 21 and 24 percent respectively. Phosphorus in shoot decreased 6 and 8 percent on the application of NS solution: @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60mM phosphorus decreased in shoot 13 and 15 percent respectively. Phosphorus in root decreased 6 and 11 percent on the application of NS solution @ 30 and 60 mM respectively. On the application of AS solution @ 30 and 60 mM phosphorus decreased in root 13 and 15 percent respectively (Table 2).

Sulphur and phosphorus were not applied as supplementary nutrients, but as per standard recipe. In both types of salt stress, the concentration of sulphur and phosphorus in

shoot and root decreased. The influential parameters of neutral and alkaline salts affected sulphur and phosphorus metabolism in the growing plants. For a healthy growth of a plant it is pertinent that root and shoot system must be in co-ordination. The higher the intake of nutrients by a root system according to the growth requirements of a plant and higher is the nutrients transport and the metabolic activities. Uptake and translocation of nutrients play essential roles in plant for signal transduction, growth, and development [20]. In this study, it might be that types of salts for stress affected sulphur and phosphorus acquisition by interfering with K^+ uptake by carriers and channels as revealed by [21].

3.3 Effect of Types of Salt Stress on Na^+/K^+ in Wheat

In shoot and root, Na^+/K^+ ratio increased than the control on the application of both types of salt solutions. In addition, Na^+/K^+ ratio was higher in case of AS than NS applied solutions. The magnitude of Na^+/K^+ ratio was higher in shoot than the root (Fig. 2).

Presence of sodium ion affected osmotic property and presence of carbonates and bicarbonates affected pH of the nutrients supply from the surroundings to the roots. To increase mass and volume, turgor pressure of cell is desirable, which is also dependent on tissue water contents. This increases expansion as well as longitudinal growth of a plant tissue [22]. Under salinity, Na^+/K^+ ratio in wheat is increased [23]. Usually with application of NS, this ratio remains higher than the presence of AS. But in this study, Na^+/K^+ ratio in seedlings was higher with AS application. pH trended to be higher in the presence of AS, and may have influenced metabolic or biochemical processes of wheat.

Table 2. Effect of neutral salts (NS) and alkaline salts (AS) stresses on concentrations of sulphur and phosphorus in wheat

Salinity (mM)	Sulphur (%)		Phosphorus (%)	
	Shoot	Root	Shoot	Root
Control	0.30 a	0.29 a	0.48 a	0.46 a
NS (30)	0.29 a	0.27 b	0.45 b	0.43 b
NS (60)	0.26 b	0.25 c	0.44 b	0.41 c
AS (30)	0.24 c	0.23 d	0.42 c	0.40 c d
AS (60)	0.23 c	0.22 d	0.41 c	0.39 d

Means sharing similar letter(s) for a parameter in a column do not differ significantly at $p < 0.01$; CV ($p < 0.01$) = 7.52.98 percent




Fig. 2. Effect of neutral salts (NS) and alkaline salts (AS) on Na⁺/K⁺ ratio in shoot and root of wheat

Under genetic, this wheat variety may have inclined to accumulate more sodium ion than potassium ion in the organs. In addition, the magnitude of Na⁺/K⁺ ratio was higher in shoot than the root. Root system was genetically unable to accumulate more sodium ion than the shoot system. Wheat shows high genetic variability across the synthetic backcross lines and their parents at various growth stages under controlled as well as under salt stress field conditions [24].

4. CONCLUSIONS

The magnitude of decrease of root length, shoot height and their biomass was higher with alkaline salts than neutral salt application in the growth medium. This may be due to 7 percent lower P concentration in alkaline salts besides high Na⁺/K⁺. This study can be helpful to advise farmers to focus on types of salts in a salty soil.

COMPETING INTERESTS

Authors have declared that no competing interests exist.

REFERENCES

1. Tobe K, Li XM, Omasa K. Effects of five different salts on seed germination and seedling growth of *Haloxylon ammodendron* (Chenopodiaceae). *Seed Science Research*. 2004;14:345–353.
2. Wang H, Ahan J, Wu ZH, Shi DC, Liu B, Yang CW. Alteration of nitrogen metabolism in rice variety 'Nipponbare' induced by alkali stress. *Plant and Soil*. 2011;355:131–47.
3. Mudgal V, Madaan N, Mudgal A. Biochemical mechanisms of salt tolerance in plants: A Review. *International Journal of Botany*. 2010;6:136–143.
4. Hajjaoui H, El Ayebe N, Garrecc JP, Dendend M. Differential effects of salt stress on osmotic adjustment and solutes allocation on the basis of root and leaf tissue senescence of twosilage maize (*Zea mays* L.) varieties. *Industrial Crops and Products*. 2010;31:122–130.
5. Badr Z, Rehana A, Salim M, Safdar A, Niazi BH, Arshad A, Mahmood IA. Growth and ionic relations of *Brassica campestris* and *B. juncea* (L.) Czern & Coss. under induced salt stress. *Pakistan Journal of Agricultural Sciences*. 2006;43(3-4):103–107.
6. Yang CW, Jianaer A, Li CY, Shi DC, Wang DL. Comparison of the effects of salt-stress and alkali-stress on photosynthesis and energy storage of an alkali resistant halophyte *Chloris virgata*. *Photosynthetica*. 2008;46:273–278.
7. Shi DC, Yin SJ, Yang GH, Zhao KF. Citric acid accumulation in an alkalitolerant plant *Puccinellia tenuiflora* under alkaline stress. *Acta Botanica Sinica*. 2002;44:537–540.
8. Rengasamy P. Soil processes affecting crop production in salt-affected soils. *Functional Plant Biology*. 2010;(37):613–620.
9. Ghoulam C, Foursy A, Fares K. Effects of salt stress on growth, inorganic ions and

- proline accumulation in relation to osmotic adjustment in five sugar beet cultivars. *Environmental and Experimental Botany*. 2002;7:39–50.
10. De Lacerda CF, Cambraia J, Oliva MA, Ruiz HA, Prisco JT. Solute accumulation and distribution during shoot and leaf development in two sorghum varieties under salt stress. *Environmental and Experimental Botany*. 2003;49(2):107-120.
 11. Shi DC, Wang DL. Effects of various salt-alkaline mixed stresses on *Aneurolepidium chinense* (Trin.) Kitag. *Plant and Soil*. 2005;271:15–26.
 12. Britto DT, Kronzucker HJ. NH_4^+ toxicity in higher plants: a critical review. *Journal of Plant Physiology*. 2002;159:567-584.
 13. Hoagland DR, Arnon DI. The water culture method of growing plants without soil. *Univ. California, Berkeley College Agriculture. Circ.*1950;344.
 14. Jackson ML. *Soil Chemical Analysis*. Contable Co. Ltd. London. 1962;62.
 15. Verma BC, Swaminathan KS, Sud KS. An improved turbidimetric method for sulphur determination in plants and soils. *Talanta*. 1977;24:49-50.
 16. Statistix 8.1. *Statistical Software for Windows*, Tallahassee, Florida; 2005.
 17. Tavakkoli E, Rengasamy P, McDonald GK. High concentrations of Na^+ and Cl^- ions in soil solution have simultaneous detrimental effects on growth of faba bean under salinity stress. *Journal of Experimental Botany*. 2010;61(15):4449–4459.
 18. Badr Z, Salim M, Rehana A, Niazi BH, Mahmood IA, Ali A. Growth Responses and Ionic Relations in two *Brassica* species under water stress conditions. *Pakistan Journal of Scientific and Industrial Research*. 2008;51(1):31-35.
 19. Badr Z, Salim M, Rehana A. Role of Ca^{2+} on Growth of *Brassica campestris* L. and *B. juncea* (L.) Czern& Coss under Na^+ Stress. *Journal of Integrative Plant Biology*. 2010a;52(6):549–555.
 20. Wang D, Lv S, Jiang P, Li Y. Roles, regulation, and agricultural application of plant phosphate transporters. *Front. Plant Science*. 2017;8(817):1-14.
 21. Maathuis FJM, Amtmann A. K^+ nutrition and Na^+ toxicity: The basis of cellular K^+/Na^+ ratios. *Annals of Botany*. 1999;84: 123–133.
 22. Badr Z, Ali A, Mahmood IA, Arshadullah. M, Armaghan S, Adil MK. Potassium consumption by rice plant from different sources under salt stress. *Pakistan Journal of Scientific and Industrial Research*. 2010b;53(5):271-277.
 23. Niazi BH, Badr Z, Salim M, Athar M. Growth response, water relations and K/Na ratio in wheat under sodium and calcium interaction. *Journal of Applied Sciences and Environmental Management*. 2007; 11(1):47-50.
 24. Dadshani S, Sharma RC, Baum M, Ogbonnaya FC, Le'on J, Ballvora A. Multidimensional evaluation of response to salt stress in wheat. *PLoS One*. 2019; 14(9):1-24.

© 2019 Badar-Uz-Zaman et al.; This is an Open Access article distributed under the terms of the Creative Commons Attribution License (<http://creativecommons.org/licenses/by/4.0>), which permits unrestricted use, distribution, and reproduction in any medium, provided the original work is properly cited.

Peer-review history:

The peer review history for this paper can be accessed here:
<http://www.sdiarticle4.com/review-history/53782>